

VeyFo® Multi-Mulgat sadrži cjelovitu paletu svih važnih vitamina topljivih u vodi i mastima (osim vitamina C), kao i važne mikronutrijente poput selena. Proizvod je namijenjen upotrebi kod prasadi, svinja, teladi, janjadi, jaradi, konja, ždrebadi i kućnih ljubimaca.

Oprema:

100 ml bočica
500 ml boca

Doziranje i način primjene:

Species		Dosage
Breeding sows:	General feed supplement before parturition	1 – 3 ml daily
	After weaning off the piglets	5 – 10 ml daily
	Up to 5 days after mating	5 – 10 ml daily
Piglets:	General feed supplement	1 ml daily
	Neonates until 3 days post partum	1 ml
	Before weaning	2 ml (possibly via the drinker)
Fattening pigs:	General feed supplement	1 – 2 ml daily
Horses:	General feed supplement	5 ml daily
	5 days before to 10 days after birth	10 – 15 ml daily
	In case of increased stress	15 ml daily
Foals:	General feed supplement	1 – 2 ml daily
	Neonates immediately after birth	10 ml
	In case of increased stress	up to 4 ml daily
Dairy cows:	General feed supplement	5 – 10 ml daily
	5 to 10 days before and after insemination	15 – 20 ml daily
	In case of increased stress	
	(e.g. high daily milk yield)	5 – 10 ml daily

Calves:	General feed supplement Neonates In case of increased stress	1 – 2 ml daily 10 ml daily up to 4 ml daily
Cattle, bulls:	General feed supplement In case of increased stress	1 ml/50 kg b.w. up to 15 ml daily
Sheep, goats:	General feed supplement Dams before until approx. 10 days after lambing Before mating resp. In case of increased stress	1 – 2 ml daily 3 – 5 ml daily 4 – 5 ml daily
Newborn lambs and goat kids:		1 – 2 ml
Puppies:		2 – 5 drops per week
Mature dogs:	Small breeds: Medium-sized breeds: Large breeds:	5 – 10 drops per week, up to 20 drops in case of increased stress 20 drops, up to 30 drops per week in case of increased stress 30 drops, up to 50 drops per week in case of increased stress
Kittens:		1 – 2 drops per week
Mature cats:		2 – 5 drops per week
Poultry:	During particular demanding stages like i. e after an antibiotic therapy resp. as an extra vitamin provision	0.5 ml – 1 ml per litre drinking water (0.05 – 0.10 %) for 1 – 2 days
Pigeons:		1 drop per week, up to 2 drops in case of increased stress
Aviary birds:		1 drop every 3 rd day
Larger birds with a live weight of more than 250 g:		1 drop for every 250 g body weight every 3 rd day, up to 2 drops in case of increased stress
Reptiles:		1 drop for every 250 g body weight per week, up to 2 drops in case of increased stress
Turtles/Tortoises:		2 drops per week
Chinchillas:		1 drop every 2 nd day
Small rodents such as guinea pigs:		1 drop every 3 rd day
Rabbits:		1 drop for every 250 g body weight every 3 rd day, up to 2 drops in case of increased stress